



A STUDY OF PREVALENCE OF FISH PARASITE IN HASHENGE LAKE, TIGRAY, ETHIOPIA

Haftom Abay*

Ministry of Livestock and Fisheries, Export Abattoir Inspection and Certification Directorate Adissababa, Mekelle University, Ethiopia.

Received date: 13 November 2017

Revised date: 04 December 2017

Accepted date: 25 December 2017

Corresponding author: Haftom Abay

Ministry of Livestock and Fisheries, Export Abattoir Inspection and Certification Directorate Adissababa Ethiopia.

Email ID: abayhagos24@gmail.com.

ABSTRACT

This study was conducted at Lake Hashenge, Tigray, Ethiopia from the period of November to April 2009 to identify parasites of fish. The objective of this study was to identify parasite of fish at Lake Hashenge to determine the prevalence of parasitic disease of fish, to have baseline information fish parasite in The lake Hashenge. A total of 442 different species of were sampled. These include 217 of Mirror carp and 225 of *Oreochromis niloticus*. The fish were thoroughly examined both externally and internally for the presence of parasites. The major parasites identified were *CLinostomum* species from 96(44.2%) of Mirror carp and 64(28.4%) of *Oreochromis niloticus*. *Clinostomum* species were recovered from the faeces of the fish (gastrointestinal tract). The nematode genera of *capillaria* identified from the GIT of 13(6%) Mirror carp and 14(6.2%) *Oreochromis niloticus*. The genera of *Acanthocephalus* were recovered from the GIT of 29(13.4%) of mirror carp and 25(11.1%) *Oreochromis niloticus*. The adult of the genera *camllanus* was recovered from 20(9.2%) of Mirror carp and 27(12%) of *Oreochromis niloticus*. The $P > 0.05$ in the comparison between the fish species, sex, with genera of the parasite but the $P < 0.05$ in the comparison between sample source and occurrence of the parasites.

KEYWORDS: Fish disease, Lake Hashenge fish parasites.

1. INTRODUCTION

Fish are the most diverse group of vertebrates occupying a variety of marine and fresh water habitat; they are cold-blooded or poikilothermic animals. Their body temperature varying passively in accordance with the temperature of the surrounding water. Although fish as a group are tolerant of a wide range of temperatures. From just below zero degree centigrade up to 45 degrees centigrade. Individual species generally have a preferred optimum as well as a more restricted temperature range. Fish has enormous importance to human beings. They provide food for millions of people, in addition fish are important in the balance of nature, and they eat plants and animals and in turn become food for plants and animals. That's way fish helping in the balance of the total number of plants and animals on the earth (Fronson, 1916).

Catch fish industries has declined significantly and decline is due to a series of factor which include over fishing, loss of fish habitans and environmental pollution. Disease out breaks (infectious and non

infectious) resulting high mortalities occur more often when fish are held under relatively crowded and confined conditions. Also mass mortality of healthy fish may occur even under good environmental conditions when an infectious agent is accidentally introduced in to the culture system (Freders, 1195).

Water is the most important constituent in which fish are stocked, fed and raised and into which their waste products are voided, changes in the environment affect fish more rapidly and profoundly than terrestrial animals. The adaptation of the aquatic habit has many implications for the structure and physiology of fish. The higher density of water when compared with air makes the streamlining and shaping of the body on important prerequisite for successful aquatic life. Each species of fish has preferred ranges for the various parameters of water quality, such as temperature, dissolved oxygen and salinity and ideally the fish should operate at the optimum levels of each parameter to achieve fast growth and efficient performance (Nagasawa, 1979).

Ethiopia possesses many lakes which are important for the habitat of fish. In Ethiopia they are over 100 species of fish which are identified. Ethiopia has many lakes, but their contribution to the economy of the country is insignificant. The main factors which influence health, growth and survival of fish are water quality, environmental changes, manmade problems and presence of competitive aquatic host inside the habitat (Edward, 1978).

Lakes are exposed to many pollutants including untreated sewage, agricultural and industrial wastes which enhance the concentration of heavy metals and compromise the health status of fish and its growth at regular period. In developing countries farmers found around the lake send their animals (sheep, cattle, horse, and donkey) towards the lake for the grazing and drinking purpose. These animals defecate and urinate at the shore or bank of the lake which serves as the main source of pathogen for fish health. In addition to this farmers wash their cloth at the border of the lake by using detergents which are harmful for growth and survival of fish (Donald, 1993).

Fish have a very full complement of disease like all animals with which to contend and many of these are due to external agencies which others arise internally. From external source such as viruses, bacteria, fungi and parasites are known to affect fish while internally they suffer from almost all the common organic and degenerative disorders such as neoplastic conditions in water, from predators and the capturing devices of fishermen even if they are not killed by disease or disordered (Lagler et al, 1971).

Fish parasites can spoil the appearance of fish that result in consumer rejection. In our country there is no experience of farming of fish in inland waters of the country at this stage and no knowledge of disease in fish to be formed in these waters along with the growing interest in the development of disease as one of the major detrimental factors in culturing of fish. At present prediction of potential health hazards in largely extrapolated from data available on fish disease and infections in natural and semi natural habitats (Perna, 1980).

From the parasite of fish Nematode, Trematode and Acanthocephala are the most economical important helminthes of parasite of fishes of the world (Hafesteinsson and Rizvi, 1987). Most adult nematodes are found in the intestine of fish but the larval stage in the flesh and viscera which causes disease and economic problems. It is also the larval stages which are infective to humans (Margolis, 1977). Which have the greatest impact on consumer acceptance of fish as a source of protein (Hagasawa, 1989).

Acanthocephalans are endoparasitic worms and about one third of the species are found as adults in the

intestine of fishes. Acanthocephalans require vertebrate animals for definitive host and arthropods for intermediate hosts. Isopods, amphipods and ostracods are the usual intermediate hosts for aquatic species. Infection occurs when a definitive host consumes the infective stage contained in an arthropod or in a paratenic host worms are typically recruited in to fish populations during the spring with maturation egg production and transmission to intermediate hosts in the summer and early autumn. Adult worms usually live about one (Hagasawa, 1989).

Mucosal tissue is damaged at the attachment site, in fibroplasias that may extend through the sub mucosa and into the muscularis. Occasionally perforation of the gut wall occurs, the mucosal epithelium frequently is compressed or eroded along the trunk of the worm and the tips of the villi may be absent. Destruction of intestinal villi and necrotic, degenerative changes in mucosal epithelium almost certainly will reduce the absorptive efficiency of the fish intestine (Bullock, 1963).

Acanthocephalans lack an alimentary tract and hence uptake of nutrients derived both from leakage of host tissues and from dietary contents in the intestinal lumen of the host is through the tegument. Hydrolytic enzyme activity at the tegumental surface probably assists in obtaining nutrients and in rapid penetration by the worm. Some acanthocephalans penetrate deeply and induce formation of a nodule that extends in to the coelom of the host. Such nodules are extensively vascularized. Increased leakage of proteins from the blood in to the nodules ensures steady supply nutrients for parasite (Chubb, 1982). The consequences of acanthocephalans induced reductions in energy efficiency and altered basal metabolism of host are likely to be focused more sharply with increasing emphasis on aquaculture (Bristol, 1984). Digenetic trematode is one of the major fish parasitic plathyhelminthes. Digeneans require more than one host to complete their life cycle and their adult stage is parasitic in vertebrates. All major groups of vertebrates serve as hosts for adult digeneans. Aside from being hosts to adult digeneans, fish may be also be infected by the metacercarial larval stage. With one exception, mollusks; members of the genus *Aporocneotyle* complete their larval development in polychaete annelids (Donges, 1974).

Metacercarial infection in fish is the main source of disease with subsequent economic loss. Metacercariae may affect growth and survival, or disfigure fish so that they lose their market value as a food or ornamental product (Perna, 1980). Digenean infection of fish required by direct cercariae penetration are linked to seasonal occurrences of snails and their levels of infection (Brooks, 1985).

Adult stage digeneans have a dorsoventrally flattened oval body with a smooth, spiny or corrugated surface, a

sucker around the antero-ventral mouth and an additional ventral sucker or acetabulum. Both suckers used for attachment and locomotion. The digestive system consists of a pharynx connected to the mouth opening a short esophagus and two blinded intestinal caeca. The excretory system includes flame cells connected to a duct network which is connected to a posterior opening bladder. Most trematodes are hermaphroditic (with cross-insemination). The male system consists of testes, vasa effrentia and vas deferens, and a cirrus pouch containing a seminal vesicle, ducts, ejaculatorius and prostate glands. The female system is comprised of an ovary and paired vitellaria, an oviduct, vitelline ducts (Dollfusetal.1974).

So far, very few diseases have been described from fish of Ethiopian waters. Moreover there is no experience in farming of fish in inland waters of the country at this stage and no knowledge of disease in fish to be farmed in these waters (Eshetu, 2000).

Therefore, the objective of the current study is:

- To identify the parasite of fish in lake Hashenge.
- To determine the prevalence of parasite disease of fish in Hashenge.
- To have base line information about fish parasites in the lake.

2. MATERIALS AND STUDY METHODOLOGY

2.1 The study area

The study was conducted in Hashenge lake which is found in southern part of Tigray, Ethiopia, particularly in the woreda Ofla near to the town of Korem, which is 150 Km faraway from Mekelle, the capital city of Tigray region. The lake is oval in landscape and covering an area of 20 Km² and the capacity the lake to contain water is supposed to be 22.6X10⁸ m³. Hashenge Lake is the largest lake from Tigray region and has an average depth of 21 meter. The woreda Ofla has an annual rain fall of 800-900 mm and temperature in the range of 15-16 c0 and is located at longitude of 390 (east) and latitude 12031' (north). It has an altitude of 2450m above sea level. lake is natural and used as water source for both human being and animals, irrigation of agricultural land and nowadays used for fish farming system however before nine years the lake was not used for farming of fish because it did not have its own original fish species, but the agricultural bureau of Tigray region had been introduced fish species from other lakes of Ethiopia.

2.2. Study animal

The study population was different fish species (Oreochromis niloticus or Nile tilapia and Mirror carp) found at Lake Hashenge.

2.3 Study design

A cross sectional study was conducted from November 2008 to April 2009 at Hashenge lake to determine the types of fish species and the prevalence of parasitic

infestation or infection of fish .The desired sample size was calculated using the formula given by Thrusfield, 1955. 95% confidence interval, 5% desired absolute precision was considered as statically significant at 50% expected prevalence and by which 384 fish were sampled. But 15% was added to increase the precision of the estimated prevalence of the disease. Therefore, 442 fish were selected (formula step given below) using random systematic sampling method.

$$N = \frac{1.96^2 \times P_{ex} (1 - P_{ex})}{d^2}$$

Where

N= required sample size; p_{ex} =expected prevalence

d² =desired absolute precision

2.4. Questionnaire survey

Questionnaire format was developed to gather information regarding the dominant fish species in the lake, the origin of fish where they came from and to assess for any incidence of loss of the lake population (fish species) and administered to 60 persons that include fishermen (n=20, 33.3%), restaurant workers (n =15, 25%) and farmers (n=25, 41.7%) and all of them were living at the villages vicinity to the lake.

2.5. Study methodology

2.5.1 Sample collection

A total of 442 fish were sampled and of which 217 Mirror carp and 225 Oreochromis niloticus species of fish were examined. All the fish were caught using gillnets with mesh size ranging from 60 to 120 mm that was used for the fishing practice at the lake Hashenge. All the specimens taken from fish were preserved in 10% formalin until it reaches to the place of laboratory examination, Mekelle University College of veterinary medicine parasitological laboratory. During the process of sampling, fish species, sex and site of sample taken was identified and recorded.

2.5.2 Sampling procedure

The fish were examined thoroughly both internally and externally after it has been taken out by the fisherman from the lake, scraping were taken from the skin and fin ,preserved in 10% formalin and sample was taken from the gill and gastro-intestinal tract. An incision was made from the ventrum to the head then faeces have been collected from the small and large intestine.

2.5.3 Laboratory examinations

Skin scraping was taken from randomly selected fish type particularly from the route of the fin. The skin scraping has been preserved in 10% formalin and taken to the laboratory. In the laboratory scrapings were dissolved with 10% KOH and smear was examined using the 10x magnification power of the light microscope. Any sample which was not showing the typical intact parasites or whole egg was considered to be negative.

Fecal direct to microscopy procedure was conducted to see parasitic eggs, larvae and adult parasites from the specific samples from each fish species. Fecal sedimentation and floatation techniques have been carried out to identify parasite egg and adults. Gill samples were also examined to assess external parasitism.

2.6 Data analysis

The obtained data from questionnaire survey and laboratory findings were summarized and then analyzed using simple mathematical calculations and SPSS version 10 analyzing software to calculate the prevalence rate of the parasitic infestation in the study fish species and chi-square test was calculated to test association between the fish species, sex and site or sample source with the occurrence of parasitic infestation.

3. RESULTS

3.1 Parasitological survey

The species *Clinostomum* were the most prevalent parasites affecting both species of Fish at Hashenge Lake. Trematode parasites were recovered from 96(44.2%) of Mirror carp and 64(28.4%) of *Oreochromis niloticus* from the gastro intestinal tract of the fish. The eggs of genus *Acanthocephalans* were recovered from 29 (13.4%) of M.carp and 25(11.1%) of *O.niloticus* of fish from its gastro intestinal tract. The genera *capillaria* were found in 13(6%) Mirror carp and 14(6.2%) of *Oreochromis niloticus*. The adult stage of *Camallans* was found in 20(9.2%) of Mirror carp and 27(12%) of *Oreochromis niloticus*.

Table 1: Comparison between species of fish and parasitic prevalence.

Fish species	Number of examined fish	Number of fish affected	%
Mirror carp	217(49.1%)	158	72.8
O.niloticus	225(50.9%)	130	57.78
Total	442(100%)	288	65.11

Table 1 Showed that the overall prevalence of the fish parasite was 65.11% in the study area and 72.8 % was for Mirror carp and 57.78% for the *Oreochromis niloticus*.

from 96(44.2%) of Mirror carp and 64(28.4%) of *Oreochromis niloticus* and the minimum number of parasites belonging to the genera of *Capillaria* recovered from 13(6%) of Mirror carp and 14(6.2%) of *Oreochromis niloticus*.

Table 2 Indicate the maximum number of parasites found was belonging to the genera of *Clinostomum* recovered

Table 2: Prevalence of fish species versus parasites.

Fish species	Parasite genera				%
	<i>Clinostomum</i>	<i>Capillaria</i>	<i>Camallanus</i>	<i>Acanthocephalans</i>	
Mirror carp(n=217)	96	13	20	29	72.8
O.niloticus (n=225)	64	14	27	25	57.9
Total	160	27	47	54	65.1

$X^2 = 16.052, P(z) < 0.05, df = 4$

Table 3: Prevalence of sex versus parasite.

Sex	Parasite genera				%
	<i>Clinostomum</i>	<i>Capillaria</i>	<i>Camallanus</i>	<i>Acanthocephalus</i>	
Female (n=188)	64 (34%)	13(7)	17(9%)	18(9%)	59.57
Male (n=154)	96(38%)	14(5.5%)	30(12%)	36(14%)	69.3
Total (n=442)	160(36%)	27(6%)	47 (11%)	54 (12%)	65

$X^2 = 6.345, P (z) > 0.05, df = 4$

As seen from table 3, there was no any significant ($X^2 = 6.345, P (z) > 0.05$) variation among the sex categories of the prevalence rate of the parasite.

Table 4: Infection prevalence variation among site (sample source).

Sex	Parasite genera				%
	<i>Clinostomum</i>	<i>Capillaria</i>	<i>Camallanus</i>	<i>Acanthocephalus</i>	
Skin/fin (n=50)	0	0	0	0	0
GIT (n=328)	160	27	41	54	63
Gill (n=64)	0	0	0	0	0

$X^2 = 287.29, P (z) < 0.05, df = 8$

As we can see from table 4, there was a significant variation among the site versus parasite categories. Most of the parasite egg genera were found from the gastro-

intestinal tract (63.8) while no parasite was found from the samples taken from the gill, skin and fin.

3.2 Questionnaire survey

Table 5: Questionnaire survey.

Respondent	Question item					
	Type of fish species		Any disease outbreak		Origin of fish	
	O.N & M	Barbus	Yes	No	Haik	Tana
F.M(N=20)	18(90%)	2(10%)	5(25%)	15(75%)	8(40%)	12(60%)
R.W(N=15)	14(93%)	1(6.7%)	5(33.33%)	10(66.2%)	10(66.2%)	5(37.8%)
Farmers(n=25)	0	0	15(60%)	10(40%)	15(60%)	10(40%)
Total(n=60)	53.3%	5%	41.6%	58.33%	55%	45%

NB: F.M =fishermen, R.W= Restaurant workers

From the table 5, the most dominant fish species in the lake were Mirror carp and *Oreochromis niloticus* (53.33%) according to the questionnaire survey of the respondent indicated while 5% of the respondent said barbus species.

Hashenge Lake was a dead sea which has not its original fish but according to the Questionnaire survey the fish species present in lake were brought from Lake Haik, 272 Km far from Mekelle, and this was supported by 55% of the respondents while 28.33% of the respondent said from Lake Tana. The questionnaire survey indicated that there was an outbreak in 1993 E.C. all the fish have been found dead but the cause of mortality was not known and this previous history of the occurrence of death was supported by 41.67% of the respondents. On the other hand, 58.33% of the respondents replied that they did not have any information about such phenomenon.

4. DISCUSSION

The helminthes egg parasites belonging to the genera *Clinostomum*, *Capillaria* and *Acanthocephalus* and the adult stage of the genera *Camallans* was found in the study fish species. The study was conducted in a total of 442 fishes from Lake among which 217(49.1%) were Mirror carp and 225 (50.9%) were *Oreochromis niloticus*. From the total examined fishes, 290(65.6%) was found to be positive for the genera of parasites. Comparisons were made among each other of the different variables.

The most prevalent egg of trematode was the egg of the genus *Clinostomum* which was recovered from the gastrointestinal of *Oreochromis niloticus* (28.4%) and Mirror carp (44.2%). The genus *Clinostomum* was the most abundant parasites found in the *Oreochromis niloticus* and Mirror carp. This finding of study was consistent the finding recorded by Teferra wondum (1990), with a prevalence rate of 74.3% in *Oreochromis niloticus* and 42% in the Mirror carp at Lake Tana. These findings were slightly higher than the finding of the current study in which prevalence rate of 28.4% for *Oreochromis niloticus* and 44.2% for Mirror carp was

identified. This difference in prevalence rate may be due to management of the water quality, feeding habits and overcrowding of fishes, presence of aquatic birds that contribute to the exposure of fish to the parasite (Anderson, 1992).

The nematode of *Camallanus* was found from 20(9.2%) of Mirror carp and 27(12%) of *Oreochromis niloticus* in the present study at the study lake. The result of current study agrees with the Fujita (1927) and is reported from fresh water fishes in Japan and is reported from many hosts from many countries. Nematode usually considered as the most economically important parasite of fishes of the world (Hafsteinsson and Rizvi, 1987).

The genus *Acanthocephalus* was also found from Mirror carp (13.4%) and *Oreochromis niloticus* (11.1%). The genus *Capillaria* was found from 13(6%) of Mirror carp and 14 (6.2%) of *Oreochromis niloticus*.

The questionnaire survey revealed that the total respondents participated in the study were provided with query type of disease outbreak in the lake previously, about the origin and type of fish species and dominant fish species in the lake at present. Since fish is the main source of protein as food for human being great consideration should be given to the health status of the fish but in developing countries like Ethiopia, Fish farming is not greatly practiced (Bullock, 1963).

Farmers who are dweller in the surrounding of Lake Hashenge do not consume fish meat. From the total respondents interviewed; 41.6% of them stated that there was an outbreak of fish disease in the lake while 58.33% of the respondents replied that they did not have any information about such situation of occurrence of disease outbreak in the lake. The most dominant fish species in the lake were Mirror carp and *Oreochromis niloticus*.

5. CONCLUSION AND RECOMMENDATIONS

Fisheries have a very full complement of disease like all animals. The disease may be due to internal and external sources. The present work which was conducted in Hashenge Lake, Tigray region, revealed that the

prevalence of the parasites that affect fish were high, this may be due to lack of management of the lake and shortage of education about the importance of fish as source of food. The contribution of the farmers to the lake management was very poor and no attention was given to it at all.

Based on the above conclusion the following recommendation was forwarded:

- Creation of awareness about the importance of fish as source of food for society
- Good water and environment management of the lake
- Fish species which can be best fit to the climatic condition of the study area and water ecosystem should be selected and introduced to lake.
- The governmental should be involved in the overall management of the water body and financially support for the successful start of fish production from the lake as to the desired level.
- Training of fishery professionals about the occurrence of fish diseases should be practiced and further research should be conducted in order to increase the productivity of the lake.

ACKNOWLEDGEMENTS

My special thanks go to my advisers Dr. sisay weldegebriel and Dr. Etsay kebede. Also I would like to express my special thanks to Dr. shewit kalyou (DVM, MSC, Assis.prof) for his help in analyzing my data.

REFERENCES

1. Anderson, R.C. Nematode parasites of vertebrates and their development transmission. CAB international, 1992; 578.
2. Balarin, J. D. National reviews for aquaculture development in Africa: Ethiopia. FAD fisheries circular No., 1986; 770(9): 120.
3. Bristol, J. R., Mayberry, L. F. Huber, D. and Ehrich, 1. endoparasite fauna of Trout in the plitvice lakes National park. Veterinary Arhiv, 1984; 54: 5-11.
4. Brooks, D.R.O. Grady R.T and Glen D. R. phylogenetic analysis of the Digenea (plathyhelminthes: cercomeria) with comment on their adaptive radiation. Canadian Journal of Zoology, 1985; 63: 441-443.
5. Bullock, W.L. International history of some salmoid fishes with particular reference to the histopathology of acanthocephalan infections. Journal of morphology, 1963; 112: 23-24.
6. Chubb J.C. Seasonal occurrence of helminthes in fresh water fishes part IV. Adult cestoda, Nematode and Acanthocephala. Advances in parasitology, 1982; 20: 1-292.
7. Dollfus, R.P. Miscellanea helminthological maroccana xiii-xii. Archives de L institute pasture du maroc, 1954; 4: 583-656.
8. Donald R.swift, Aquaculture Training manual second edition, USA, 1993.
9. Donges, J. The life history of clinostomum heterostomum (Rudlph, 1809) (trmatoda: clinostomatidae). International Journal for parasitology, 1974; 4: 79-90.
10. Edwards, D.J. Salmon and trout farming in Norway. Fishing news books, 1978.
11. Eshetu Yimer. Preliminary survey of parasites and bacterial pathogens of fish at Lake Ziway, Ethiop.j.sci (sinet), 2000; 23(1): 25-33.
12. Fish diseases and disorders, protozoan and metazoan infections, P. T. K. Woo, 1995; 1.
13. Frederick, V.T. Fish Aquaculture and experiments, federal research center for fisheries, institute for coastal and inland fisheries, Hamburg (Ahrensburg Federal Republic of Germany), 1985.
14. Fonson, M.A, Standard methods for the examination of water and waste water method Washington, D.C: American public health association, 1976.
15. Hafsteinsson, H and Rizus. S, A review of the seal worm problem: Biology, implications and solutions. Journal of food protection, 1987; 50: 70-84.
16. Jeane and Ely, Introduction to Aquaculture, Mattew Landau, Stocken state collage, 1989.
17. Lee, I.S. Commercial cat fish farming Danville, Interstate printers and publishers, 1981.
18. Mozgovai, A.A. (Ascaridata of animals and man and the diseases caused by them) in: skrjabin, K.I.ced.) Principles of Nematology volume II parts 1 and 2. Izdatelstvo Akademinauk SSR, 1953.
19. Nagasawa. K notes on parasites of aquatic organisms 4. Visceral adhesions of high-seas sockeye salmon caused by the nematode philonema oncorhynchi. Aqua biology, 1989; 11: 320-321.
20. Paggi,L, Nascetti,G, Cianchi R, Orecchia, P, Mattiucci, S.D.Amello, S.Beralnd, B.Brattey, J.W and Bullini, L. Genetic evidence for three species with in pseudoterranova decipiens (Nematode , Ascridea) in the north Atlantic and Norwegain and Barents seas. International Journal for parasitology, 1991; 21: 195-221.
21. Paperna, I. Parasites, infections and diseases of fish in Africa CIFA, Tech. pap, 1980; 7: 216.
22. Paperna, I. Parasites infections and disease of fish in Africa. Food and Agricultural organization UN, CIFA Technical paper, 1980; 7.
23. Pillay, Aquaculture principles and practices, former program director aquaculture development and coordination program. Food and Agriculture organization of the United Nations Rome, Italy, 1993.
24. Senas. Desilva, 1992. Reservoir fisheries of Asia, proceedings of the 2nd Asian reservoir fisheries workshop held in Hangzhou, People's Republic of china, 15-19 October 1990.