

QUAGMIRE OF CONGENITAL CARDIAC DEFECTS IN ASSOCIATION WITH DEXTROCARDIA AND SITUS INVERSUS: RARE CASE REPORT AND REVIEW OF LITERATURE

Akhil Mehrotra^{*1}, Mohammad Shaban², Faiz Illahi Siddiqui² and Anjali Chouhan²

¹Chief, Pediatric and Adult Cardiology, Prakash Heart Station, Nirala Nagar, Lucknow, UP, India.

²Cardiac Technician, Prakash Heart Station, Nirala Nagar, Lucknow, UP, India.

Received date: 08 February 2025

Revised date: 28 February 2025

Published date: 18 March 2025



*Corresponding Author: Akhil Mehrotra

Chief, Pediatric and Adult Cardiology, Prakash Heart Station, Nirala Nagar, Lucknow, UP, India.


ABSTRACT

Situs inversus with dextrocardia is the complete reversal of the position of the abdominal and thoracic viscera. It may remain asymptomatic or could be detected early in infancy or childhood, when associated with other congenital malformations especially of the cardiovascular system. Situs inversus with dextrocardia usually exists without co-existing congenital heart disease. We report the case of a nine year female child who presented with complaints of recurrent cough and bluish discoloration of the lips. Examination revealed central cyanosis with grade 3 digital clubbing and a right-sided apical impulse. Chest radiograph showed a right sided heart and Transthoracic Echocardiography (TTE) confirmed multiple cardiac anomalies: Dextrocardia, Situs inversus, Complete ario-ventricular canal defect (CAVCD), Double outlet right ventricle (DORV) with Pulmonary Stenosis (PS).


KEYWORDS: Dextrocardia, Situs inversus, Double outlet right ventricle, Pulmonary valvular stenosis, Complete atrio-ventricular canal defect, CAVCD, Hypoplastic mitral valve annulus.

INTRODUCTION

Dextrocardia is a congenital anomaly in which the heart is situated on the right side of the body^[1] (Figure 1).



(A)



(B)

Figure 1: (A) & (B) Diagrammatic Illustration of Dextrocardia with Situs Inversus.

Situs inversus is a condition in which the major visceral organs are mirrored from their normal positions, that is the morphological left atrium is on the right while the morphological right atrium, the liver and gall bladder are on the left.^[2] This is otherwise known as Situs inversus totalis or mirror image dextrocardia (Figure 2), which is

a rare condition with a prevalence of 1:10,000 in some populations.^[3,4] It is generally asymptomatic except when associated with congenital heart defects which is seen in 5 to 10% of cases.^[3,4]




Figure 2: Cadaveric specimen of Dextrocardia with Situs Inversus: The apex of the heart is pointing to the right (Dextrocardia) in the thoracic cavity. The liver is located on the left side and stomach is on the right side.

Classification of congenital dextrocardia^[5]

Type I dextrocardia (mirror-image dextrocardia): In Type I dextrocardia, the anatomic right atrium and right ventricle are situated to the left of and anterior to the corresponding systemic chambers. A mirror-image arrangement of the cardiac chambers is present since the frontal relationship of the chambers is reversed, yet, the anteroposterior arrangement is normal.

Type II dextrocardia (dextroversion complex): In Type II dextrocardia, the relations of the cardiac chambers are normal; the right atrium and right ventricle are situated to the right of and posterior to the corresponding systemic chambers. In complete dextroversion the cardiac apex is situated anteriorly and to the right; in incomplete dextroversion or mesoversion, it is located in the substernal region and the longitudinal axis of the heart is parallel to the midsagittal axis of the chest.

Type III dextrocardia (mixed dextrocardia): In Type III dextrocardia is characterized by inversion of the atria alone or of the ventricles alone. The arrangement of the cardiac chambers is therefore in part similar to that of Type I and of Type II dextrocardia.

Type IV dextrocardia (congenital dextroposition): In Type IV dextrocardia, the heart is in the midchest, but the arrangement of the chambers is normal and the cardiac apex is still directed to the left and anteriorly. Types I, II, III, and IV represent the intrinsic group of dextrocardia, since the cardiac heterotaxy is caused by a developmental anomaly of the primitive heart tube.

Type V dextrocardia (congenital extrinsic): In Type V dextrocardia, the abnormal position of the heart is due to its displacement by congenital anomalies of the lungs, diaphragm, or chest cage. The rightward displacement may be in the frontal plane only (simple dextroposition) or may occur in both the frontal and horizontal planes (dextroposition with pivotal rotation).

The incidence of cardiac anomalies in the intrinsic group was 79 per cent, and in the extrinsic group 24 per cent.^[5] These anomalies were, as a rule, multiple and severe and together formed specific anatomic syndromes usually characterized clinically by dextrocardia, cyanosis, and diminished pulmonary blood flow.^[5] The most common entities encountered were tetralogy, single ventricle with pulmonary stenosis, and tricuspid atresia or stenosis.^[5]

Here we are reporting a case of a nine year old female child presenting with dextrocardia, situs inversus accompanied with conundrum of congenital heart anomalies.

CASE REPORT

A nine year old female child was referred to us for comprehensive color echocardiography and opinion regarding management of cyanotic congenital heart disease.

The parents provided the history of cyanosis since birth, shortness of breath and failure to thrive. On clinical examination, the child was thin built, very active and lively. Pectus excavatum was noticed along with prominent chest wall. Central cyanosis was obvious over her lips, fingers tips and toes. Conspicuous clubbing was also noticed over nails of fingers and toes (Figure 3).



(A)



(B)




Figure 3: Facial appearance and other clinical features in our index patient. (A) Facial appearance; (B) Chest wall abnormalities- Pectus excavatum was observed alongwith prominent chest wall; left > right; (C) Clubbing and cyanosis was detected in all the fingers; (D) Clubbing and cyanosis was noticed in all the toes.

The child's weight was 13 kg, height was 168 cm, BP was 90/60 mmHg, HR was 105/min, respiratory rate was 16/min and SPO₂ was 65% at room air. All the peripheral pulses were normally palpable without any radio-femoral delay. Cardiovascular examination revealed apical impulse on the right side of chest, in the 5th intercostal space, just medial to the mid-clavicular

line. A grade 3/6 ejection systolic murmur was best heard in the right IInd intercostal space. P₂ component was soft. No clicks or gallop sound were heard.

Xray chest (PA) view demonstrated dextrocardia with reduced pulmonary blood flow (Figure 4).




Figure 4. X-ray chest (PA) view- Dextrocardia is recognized. The pulmonary blood flow is decreased.

Resting ECG showed the following features (Figure 5):

- Normal sinus rhythm with a ventricular rate of 96/min,
- T wave inversion in L1, AVL, V2- V6,
- Tall R waves in V1 (consistent with right ventricular hypertrophy),
- lack of normal R wave progression in precordial leads (compatible with dextrocardia).



Figure 5: Resting ECG demonstrated: Normal sinus rhythm with a ventricular rate of 96/ min, T wave inversion in L1, AVL, V2- V6 , Tall R waves in V1 (consistent with right ventricular hypertrophy), lack of normal R wave progression in precordial leads (compatible with dextrocardia).

Transthoracic Echocardiography

All echocardiography evaluations were performed by the author, using My Lab X7 4D XStrain echocardiography machine, Esaote, Italy. The images were acquired using a adult probe equipped with harmonic variable frequency electronic single crystal array transducer while the subject was lying in supine and left lateral decubitus positions.

Conventional M-mode, two-dimensional, pulse wave doppler (PWD), continuous wave doppler (CWD) and

sequential segmental transthoracic echocardiography was performed in the classical subcostal, parasternal long axis (LX), parasternal short axis (SX), 4-Chamber (4CH), 5-Chamber (5CH) and suprasternal views.

M-mode Echocardiography

M-mode echocardiography of left and right ventricle was achieved for estimating various ventricular parameters (Table 1, Figure 6).

Table 1: Calculations of M-mode echocardiography.

| Measurements | LV | RV |
|--------------|---------|---------|
| IVS d | 6.4 mm | 7.2 mm |
| ID d | 12.1 mm | 21.4 mm |
| PW d | 4.3 mm | 8.3 mm |
| IVS s | 7.2 mm | 7.0 mm |
| ID s | 8.0 mm | 16.6 mm |
| PW s | 5.9 mm | 8.3 mm |
| EF | 67 % | 48 % |
| %FS | 33 % | 22 % |
| EDV | 3.4 ml | 15.2 ml |
| ESV | 1.1 ml | 7.9 ml |
| SV | 2.3 ml | 7.3 ml |
| Mass | 9 g | 34 g |

IVS, interventricular septum, ID, internal dimension; PW, posterior wall, d, diastole; s, systole; FS, fractional shortening; EDV, end-diastolic volume; ESV, end systolic volume; SV, stroke volume; EF, ejection fraction.




Figure 6: M-mode Echocardiography. (A) LV volumetric estimation; (B) RV volumetric estimation.

Summary of M-mode echocardiography

M-mode echocardiography depicted dilated RV with mildly reduced RVEF (48%). On the contrary the LV was hypoplastic with normal LV systolic function LVEF (67%).

2Dimensional-Transthoracic Echocardiography

2Dimensional transthoracic echocardiography (TTE) was conducted in explicit detail and it demonstrated the following features (Figures 7-18):

- Dextrocardia
- Situs inversus

- Concordant L – Bulboventricular Loop, AV Concordance, VA Concordance
- L-loop ventricles
- Morphological RV is to the left of morphological LV
- Confluent pulmonary arteries
- Double outlet right ventricle (DORV)
- Both the great arteries are arising from the morphological right ventricle.
- A subaortic conus was recognised
- Malposition of great arteries

Aorta is anterior and to the right
 Pulmonary artery is posterior and to the left

- Complete atrio - ventricular canal defect (CAVCD) - Rasteilli Type B

Common AV valve (CAAV) orifice
 Common AV valve with separate MV (Right AV valve) and TV (Left AV valve) orifices
 MV orifice was hypoplastic and TV leaflets were large and thickened.
 Atrial septal defect (large)
 Size: 9.8 mm.

Ostium primum type.
 Lt. to Rt. shunt.
 Ventricular septal defect (large)
 Size 15.9 mm.
 Inlet type
 Lt. to Rt. shunt.
 Rasteilli Type B

The chordae of left AV valve (Tricuspid valve) were attached to the lateral wall of RV and RV apex. MV chordae were attached to ventricular septal crest and lateral wall of LV.

- Left AV valve (TV) regurgitation (mild)

On color flow mapping a mild, central tricuspid regurgitation jet was identified in RA. The jet area was 2.80 sqcm.

- Pulmonary valvular stenosis (severe)
 PV domed
 Peak/mean gradient across PV = 83/64 mmHg.
 Hypoplasia of PV annulus, MPA, LPA and RPA
 PV annulus (D): 4.0 mm
 MPA (D) : 5.3 mm
 LPA (D) : 3.6 mm
 RPA (D): 3.5 mm
 - Morphological RV was dilated with mildly reduced - RVEF (48 %)
- Morphological LV was hypoplastic
 Normal LV systolic function LVEF (67%)
- Dilated RA, hypoplastic LA, LV.




Figure 7: Dextrocardia. Subcostal view identifies dextrocardia situs inversus.






Figure 8: Situs Inversus: In the subcostal view aorta is on the right side and inferior vena cava (ivc) is on the left side. Liver is also left sided; ao, aorta; ivc, inferior vena cava; sp, spine.



(A)



(B)

Figure 9: Situs Inversus. Ultrasound of abdomen reveals: (A) Spleen was on the right side; (B) Liver is on the left side; sp, spine.




Figure 10: Apical 4C view from right chest with cursor probe placed at 9 o'clock position. Morphological right ventricle was dilated and morphological LV was hypoplastic; la, left atrium; lv, left ventricle; ra, right atrium; rv, right ventricle; vs, ventricular septum; TV, tricuspid valve; as, atrial septum.




Figure 11: Double outlet right ventricle. (A) and (B) In the right LX and right apical 5C view both great arteries are arising from morphological RV. Aorta was anterior and pulmonary artery was posterior and hypoplastic. Pulmonary valve was domed. A subaortic conus was also recognized. (C) In the right apical 5C view, the diameter of hypoplastic pulmonary arteries was estimated. PV annulus, MPA, LPA and RPA diameter was 4.0 mm, 5.3 mm, 3.5 mm and 3.6 mm respectively; (D) Another image of DORV in the LX view from right chest; DORV, double outlet right ventricle, Ao, aorta, PA, pulmonary artery; C, conus; LA, left atrium; TV, tricuspid valve; RV, right ventricle; mv, mitral valve; asd, atrial septal defect; l, left pulmonary artery, r, right pulmonary artery; vs, ventricular septum.




Figure 12: Color flow mapping in apical 5C view from right chest. The classical movement of blood flow in DORV is depicted. The anterior aortic flow is non turbulent and the posterior pulmonary arterial flow is highly turbulent because of severe pulmonary valvular stenosis; la, left atrium; Iv ,left ventricle; laa, left atrial appendage; AO, aorta; PA, pulmonary artery.




Figure 13: Color flow mapping across pulmonary valve identifies pulmonary valvular stenosis with a peak/mean gradient of 83.1/ 64.8 mmHg; PS, pulmonary stenosis.




Figure 14: Complete atrio-ventricular canal defect (CAVCD). (A) Apical 4C view from right chest discerns the ostium primum ASD (**) and CAVCD (****); (B) Apical 4C view from right chest portrays common AV valve with two components: large anterior and septal leaflet of TV and bileaflet MV; mv, mitral valve; Iv, left ventricle; vs, ventricular septum; mb, moderator band; rv, right ventricle; ra, right atrium; AL, anterior leaflet of TV; SL, septal leaflet of TV; as, atrial septum; TV, tricuspid valve.




Figure 15: Complete AV canal defect apical 4C - right chest and subcostal view. In the apical 4C view from right chest delineates CAVCD with large ASD and VSD. Dilated RV and small LV with large TV leaflets and hypoplastic MV annulus was also recognized. AL, anterior leaflet; SL, septal leaflet; **, CAVCD; **, ostium primum ASD; hv, hepatic vein; ivc, inferior vena cava; la, left atrium, lv, left ventricle; ra, right atrium; rv, right ventricle; as, atrial septum; vs, ventricular septum; tv, tricuspid valve; mv, mitral valve.**




Figure 16: Left AV valve (TV) regurgitation. Apical 4C view from right chest shows mild TV regurgitation.




Figure 17: Color flow mapping across ostium primum ASD denoted as (***) and complete atrio-ventricular canal defect (CAVCD) denoted as (***) . Green arrows portray the flow of blood across ASD and CAVCD; la, left atrium; ra, right atrium; rv, right ventricle; as, atrial septum; ao, aorta; mv, mitral valve; AL, anterior leaflet of tricuspid valve.




Figure 18: Suprasternal view. (A) Left aortic arch; (B) Color flow mapping demonstrating non turbulent blood flow in the left aortic arch; as, ascending aorta; da, descending aorta.

DISCUSSION

Situs inversus totalis (SIT) is a rare congenital abnormality characterized by a mirror-image transposition of both the abdominal and the thoracic organs. This is a global defect of situs orientation, as the failure to generate normal left-right asymmetry results in a spectrum of laterality disturbances.^[6] This condition might cause difficulties during diagnostic and therapeutic procedures.^[7]

Nomenclature

The body of vertebrate species shows an external bilateral symmetry. Therefore, it is surprising that humans and all other vertebrates present an internal asymmetry: nearly all visceral organs are left–right (LR) asymmetrical in their anatomy and placement.^[6] Any failure in the normal left-right asymmetry stands within a

spectrum of laterality disturbances. Heterotaxia is the common term used for these disorders. It encompasses a wide variety of disorders which may be partial or complete. The asymmetry of unpaired organs (heart, liver, spleen, stomach, bowels, aorta, v. cava) seems to be logic and “unavoidable”. The asymmetry of paired organs seems less obvious, but they have distinct left and right forms. The lungs have major differences (three lobes vs two lobes), but even kidneys show some asymmetry. Especially the vascular differences between left and right kidneys have some technical importance, eg, for transplant surgeons.

The complete reversal of normal organ position, including both the thoracic and abdominal organs, is called situs inversus totalis. Other types of situs are situs solitus (normal) and situs ambiguus

(indeterminate), which is characterized by isomerism, heterotaxy, and multiple malformations in one or more thoracic or abdominal organs.^[8]

Risk factors - Genetic mutations

In a population-based study using multivariate analyses, six risk factors for SIT were identified: family history of heart defects, family history of noncardiac anomalies, maternal diabetes, antitussive use, paternal smoking, and low socioeconomic status.^[9] These factors might have some connection with genetic mutations, as major advances in the identification of gene mutations in animals and humans support a growing body of knowledge about the causes of laterality defects.^[10]

Situs inversus is seldom seen in identical twins^[7,11], even though in conjoined twins, the incidence of SIT is high.^[12]

Laterality is established early in development and is orchestrated by a cascade of signal molecules and genes. Importantly, the neurotransmitter serotonin (5HT) also plays a critical role in this signalling cascade that establishes laterality.^[13] Sidedness is established at the time of primitive streak formation (gastrulation) and is regulated by genes, such as Nodal and PITX2, which become restricted in their expression (Figure 19).




Figure 19: Gene expression patterns for establishing the left-right axis at the 3rd week of fetal development.

Partial reversal of left–right asymmetry is more often associated with other defects than complete asymmetry (situs inversus).^[13] The discovery of these cascades has also shed light on the cause of situs defects in conjoined twins.^[14]

Genetic deletions of KIF3-A or KIF3-B, two microtubule-dependent kinesin motor proteins, resulted in a randomization of the situs of the viscera.^[15–17] Most

cases of heterotaxy are sporadic occurrences without a recognizable cause.^[9]

Epidemiology

The incidence of all lateralization defects is approximately 1:15,000 according to the Orphanet.^[18] Heterotaxy is present in approximately 1 in 10,000 births.^[19] The incidence of situs inversus itself is reported to be 1:6500 to 1:25,000, see Table 2.

Table 2: Estimated Frequency of Left-Right axis Disorders.^[20]

| | | | | |
|------------------------|----------|----------|--------|----------|
| Heterotaxy | 1:10 000 | | | |
| Situs ambiguous | 1:10 000 | 1:15 000 | | |
| Situs inversus totalis | 1:20 000 | 1:6-8000 | 1:6500 | 1:25 000 |

Situs inversus totalis is inherited in an autosomal recessive pattern^[20] and is slightly more frequent in males: 1.5:1.^[21]

Classification of situs inversus

Situs inversus could be classified into the following:

- Situs inversus totalis, where the abdominal organs are mirror images of the normal anatomy, as well as dextrocardia.^[22]

- Situs inversus with levocardia, where the base-to-apex axis points to the left while the abdominal viscera are reversed.^[22]
- Situs ambiguus. In these patients, the liver may be midline, asplenia or polysplenia, unclear cardiac morphology, or mal-rotated gut.^[22]

Compared to the other types, situs inversus totalis is the commonest.^[23]

Associations

The common congenital cardiac anomalies associated with dextrocardia with situs inversus are atrial situs solitus (93%), discordant AV connection (44%), and discordant Ventriculo-Atrial (VA) connection (30%). Congenitally corrected Transposition of Great Arteries (TGA) occurs in less than 1% of all forms of congenital heart disease.^[24] Certain congenital anomalies such as polysplenia (left isomerism)/asplenia (right isomerism) or Kartagener's syndrome often leading to infection of the paranasal sinuses and lungs) are known to occur.^[25] About 25% of individuals with situs inversus have an association with primary ciliary dyskinesia. Situs inversus totalis with primary ciliary dyskinesia together known as Kartagener's syndrome characterized by the triad of situs inversus, chronic sinusitis, and bronchiectasis.^[26]

Situs Invesus and Organ Transplantation

Organ transplantation in people with SIT presents unique challenges due to the mirror-image reversal of their organs. Surgeons must carefully assess the changed anatomy using modern imaging investigations and have particular surgical expertise to negotiate the challenges of transplant procedures in these patients. Posttransplant care and surveillance are critical for SIT people, as they may experience a higher risk of complications that need rapid intervention.^[27,28] In circumstances where multi-organ transplants are required, coordinating numerous surgeries adds complexity to the treatment plan.^[29,30] Continued research and developments in transplant medicine are required to improve outcomes for SIT patients through innovative surgical procedures and individualized treatment plans. Addressing these concerns can assist to improve the care and treatment outcomes of SIT patients receiving organ transplants.

CONCLUSION

Dextrocardia with situs inversus is a rare congenital malformation that must be thoroughly evaluated when noticed because it may result in fatal outcomes in rare instances. There is a need for a complete and elaborate diagnostic work up of suspected cases by various imaging modalities to not be missed. Doctors should encourage routine medical examinations for their patients, which could help identify this anomaly, thereby preventing the wrong diagnosis and possibly death due to delay in management. To confirm the suspected cases of situs inversus, imaging studies should be done, such as a chest x-ray or ultrasound, and a referral may be made to

a cardiologist. Imaging studies will also rule out the possibility of random arrangement of the organs, or heterotaxy, which has a much higher risk for serious medical complications. Documenting situs inversus in an individual is essential to avoid any inadvertent clinical or surgical mishap. Our case is one of the rarest reported combinations of Dextrocardia, Situs inversus, DORV, PS, CAVCD-Rasteilli Type B with Hypoplastic MV annulus and left aortic arch.

REFERENCES

1. Wilhelm A, Holbert MJ. Situs inversus. E-medicine.
2. Capdevilla J, Belmonte JCI. Knowing left from right: The molecular basis of laterality defects. *Molecular Medicine Today*, 2000; 6: 112-118.
3. Oppido G, Napoleone CP, Martano S, Gargiulo G. Hypoplastic left heart syndrome in Situs inversus totalis. *European Journal of Cardiothoracic Surgery*, 2004; 26: 10521054.
4. Torgensen J. Genic factors in visceral asymmetry in the development and pathological changes of the lung, heart and abdominal organs. *Arch Pathol*, 1949; 47: 566-93.
5. Arcilla RA, Gasul BM. Congenital dextrocardia: Clinical, angiocardiographic, and autopsy studies on 50 patients. *The Journal of Pediatrics*, 1961; 58: 251-262.
6. Peeters H, Devriendt K. Human laterality disorders. *Eur J Med Genet*, 2006; 49: 349-362.
7. Blegen HM. Surgery in situs inversus. *Ann Surg*, 1949; 129: 244-259.
8. Aylsworth AS. Clinical aspects of defects in the determination of laterality. *Am J Med Genet*, 2001; 101: 345-355.
9. Ferencz C, Loffredo CA, Correa-Villasenor A, Wilson PD. Defects of laterality and looping. In: *Genetic and Environmental Risk Factors of Major Cardiovascular Malformations. The Baltimore-Washington Infant Study: 1981-1989*. Armonk: Futura Publishing Company. Inc, 1997; 41-58.
10. Lin AE, Ticho BS, Houde K, Westgate MN, Holmes LB. Heterotaxy: associated conditions and hospital-based prevalence in newborns. *Genet Med.*, 2000; 2: 157-172.
11. Segal NL. Situs inversus totalis in twins: a brief review and a life history/twin research: twin studies of trisomy 21; monozygotic twin concordance for bilateral coronoid hyperplasia; prenatal hormonal effects in mixed-sex non-human primate litters; insurance mandates and twinning after in vitro fertilization/news reports: first report of identical twin puppies; twins sisters turn 100; remembering an identical twin production designer; New York City Marathon Quadruplets. *Twin Res Hum Genet*, 2017; 20: 90-95.
12. Layton WM. Situs inversus in conjoined twins. *Am J Med Genet*, 1989; 34: 297.
13. Sadler TW. Langman's medical embryology. In: Yamada T, editor. *Textbook of*

- Gastroenterology. 14th ed. Philadelphia: JB Lippincott, 1995; 2.
14. Levin M, Roberts DJ, Holmes LB, Tabin C. Laterality defects in conjoined twins. *Nature*, 1996; 384: 321.
 15. Haruki T, Maeta Y, Nakamura S, et al. Advanced cancer with situs inversus totalis associated with KIF3. Complex deficiency: report of two case. *Surg Today*, 2010; 40: 162–166.
 16. Teng J, Rai T, Tanaka Y, et al. The KIF3 motor transports N-cadherin and organizes the developing neuroepithelium. *Nat Cell Biol.*, 2005; 7: 474–482.
 17. Sulik K, Dehart DB, Iangaki T, et al. Morphogenesis of the murine node and notochordal plate. *Dev Dyn.*, 1994; 201: 260–278.
 18. Orphanet. The portal for rare diseases and orphan drugs. Available from: Accessed February 125, 2022.
 19. Grimes DT, Burdine RD. Left-right patterning: breaking symmetry to asymmetric morphogenesis. *Trends Genet*, 2017; 33: 616–628.
 20. Eitler K, Bibok A, & Telkes G. Situs Inversus Totalis: A Clinical Review. *International journal of general medicine*, 2022; 15: 2437-2449.
 21. Lee SE, Kim HY, Jung SE, Lee SC, Park KW, Kim WK. Situs anomalies and gastrointestinal abnormalities. *J Pediatr Surg*, 2006; 41: 1237–1242.
 22. Mujo T, Finnegan T, Joshi J, Wilcoxon KA and Reed JC. Situs Ambiguous, Levocardia, Right Sided Stomach, Obstructing Duodenal Web, and Intestinal Nonrotation: A Case Report. *Journal of Radiology Case Reports*, 2015; 9: 16-23.
 23. Maldjian PD and Saric M. Approach to Dextrocardia in Adults. *American Journal of Roentgenology*, 2007; 188: S39-S49.
 24. Ma N, Jiang SL, Huang LJ, et al. Diagnosis of isolated dextrocardia using angiocardiography or surgery. *Chinese Medical Journal (England)*, 2004; 117: 1655–1658.
 25. Shapiro AJ, Davis SD, Ferkol T, et al. Laterality defects other than situs inversus totalis in primary ciliary dyskinesia. *Chest*, 2014; 146: 1176–1186.
 26. Dabhi AS, Chaudhari SR, Throat PB, et al. Kartagener syndrome: a triad of bronchieatasis, situsinversus and chronic sinusitis. *Journal, Indian Academy of Clinical Medicine*, 2005; 6: 241–243.
 27. Schertler T, Lardinois D, Boehm T, Weder W, Wildermuth S, Alkadhi H. Lung transplantation in Kartagener syndrome and situs inversus: potential of multidetector row computed tomography and three-dimensional postprocessing. *The Journal of Thoracic and Cardiovascular Surgery*, 2007; 134: 814–815.
 28. Alnajar A, Chen PC, Burt B, Loor G. Left pulmonary artery patch augmentation for lung transplant in a patient with situs inversus. *Texas Heart Institute Journal*, 2021; 48: e197112.
 29. Deuse T, Reitz BA. Heart-lung transplantation in situs inversus totalis. *The Annals of Thoracic Surgery*, 2009; 88: 1002–1003.
 30. Chang YL, Wei J, Chang CY, Chuang YC, Sue SH. Transplantation Proceedings. Amsterdam, The Netherlands: Elsevier; 2008. Cardiac transplantation in situs inversus: two cases reports.